

Antibacterial Activity of Honey on Cariogenic Bacteria

Fatemeh Ahmadi – Motamayel¹, Seyedeh Sare Hendi², Mohammad Yusof Alikhani³, Zahra Khamverdi⁴✉

¹Associate Professor, Dental Research Center, Department of Oral Medicine, Dental school, Hamadan University of Medical Sciences, Hamadan, Iran

²Post Graduate Student of Endodontics, Dental school, Hamadan University of Medical Sciences, Hamadan, Iran

³Associate Professor, Department of Microbiology, Medical school, Hamadan University of Medical Sciences, Hamadan, Iran

⁴Associate Professor, Dental Research Center, Department of Operative Dentistry, Dental school, Hamadan University of Medical Sciences, Hamadan, Iran

Abstract

Objective: Honey has antibacterial activity. The aim of this study was to evaluate the antibacterial activity of honey on *Streptococcus mutans* and *Lactobacillus*.

Materials and Methods: In this in vitro study, solutions containing 0%, 5%, 10%, 20%, 50% and 100% (w/v) of natural Hamadan honey were prepared. Each blood (nutrient) agar plate was then filled with dilutions of the honey. The strains of bacteria were inoculated in blood agar for 24 hours at 37°C and were adjusted according to the McFarland scale (10×10^8 cfu/ml⁻¹). All assays were repeated 10 times for each of the honey concentrations. Data were analyzed by non parametric Chi-Square test. Statistical significance was set at $\alpha=0.05$.

Results: Significant antibacterial activity was detected for honey on *Streptococcus mutans* in concentrations more than 20% and on *Lactobacillus* in 100% concentration ($P<0.05$).

Conclusion: It seems that antibacterial activity of honey could be used for prevention and reduction of dental caries.

Key Words: Honey; Antibacterial Agents; *Streptococcus Mutans* *Lactobacillus*; Dental Caries

✉ Corresponding author:
Z. Khamverdi, Dental Research Center, Department of Operative Dentistry, Dental school, Hamadan University of Medical Sciences, Hamadan, Iran

zkhamverdi@yahoo.ca

Received: 17 September 2012

Accepted: 19 December 2012

Journal of Dentistry, Tehran University of Medical Sciences, Tehran, Iran (2013; Vol. 10, No.1)

INTRODUCTION

Honey is super saturated nectar collected by bees from a wide variety of plants. Although it is very well known as a food, it is not well-recognized as a medicine [1]. The medicinal and antimicrobial activities of honey for wound healing have been introduced for approximately 4500 years, almost as long as bacteria have been known. Honey has been used

as a medicine since ancient times [2-4]. In Islamic countries, it is also based on the belief in the words regarding honey in the Holly Quran that 1900 years ago and that the Lord taught the bee to build its cells on trees and in men's habitations, then to eat all the products of earth and find with skill the spacious paths of its Lord, there issues from within their bodies a drink of varying colors, wherein is healing for

men, verily in this is a sign for those who give thought [5,6]. The composition of the honey sample depends on the composition of the nectar, from which it originates. The natural antioxidants and flavonoids exhibit a wide range of biological effects including antibacterial, anti-inflammatory, anti-allergic, anti-thrombotic and vasodilatory action [7,8]. In addition, its physical properties provide ideal moist conditions for healing and a stimulating effect on the growth of wound repair tissues [9]. Unlike other antiseptics, honey has no harmful effects on tissues because slow enzymatic production of hydrogen peroxide is one thousandth hydrogen peroxide 3% [9]. Honey contains about 181 substances. There is scientific evidence about honey in several experimental and clinical conditions; for example, it has been reported effective in the treatment of gastrointestinal disease through gastric protection against gastric lesions, healing of wounds and burns, and as an antimicrobial agent [10-13]. Honey can be kept for long periods of time without spoiling, because of its high osmotic pressure and antibacterial activity [7]. It has been noted that honey clears infection, removes malodors, reduces inflammation and pain, subsides edema and exudation, and it has healing properties by stimulating angiogenesis. Honey has the potential to be useful for periodontal therapy, prevention of infection in wounds following extraction or oral surgery, other oral infections, erosion of mucosa, radiotherapy-induced stomatitis and oral ulcers with no adverse reaction [14-18].

It has been revealed that honey has antimicrobial activity against aerobic, anaerobic, gram positive and gram negative bacteria, moulds and yeasts with unique properties because of its bacteriostatic and bactericidal effect [19]. Honey differs from other sweets because of enzymes such as invertase, diastase and glucose oxidase. Moreover, honey is effective on infections which cannot be treated with conventional drugs [18,20]. Studies have demonstrated that many bacterial and fungal patho-

gens that are sensitive to honey include *Bacillus anthracis*, *Corynebacterium diphtheriae*, *Escherichia coli*, *Haemophilus influenzae*, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Mycobacterium tuberculosis*, *Pasteurella multocida*, *Proteus* species, *Pseudomonas aeruginosa*, *Salmonella choleraesuis*, *Salmonella typhimurium*, *Serratia marcescens*, *Shigella* species, *Staphylococcus aureus*, *Streptococcus mutans*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Vibrio cholerae*, *Alcaligenes faecalis*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Mycobacterium phki*, *Staphylococcus epidermidis*, *E coli* and *Helicobacter pylori* [21-23]. Recently, studies have focused on new aspects of honey application for various therapeutic purposes, especially on its antibacterial properties. Since *Streptococcus mutans* and *Lactobacillus*, as two main cariogenic bacteria of the oral cavity, play an important role in dental caries, understanding the antibacterial effect of honey on these cariogenic bacteria is necessary.

Only limited studies have been performed about the antibacterial activity of honey against cariogenic bacteria [1]. The aim of this study was to evaluate the antibacterial activity and the ability of honey to prevent growth of these bacteria (*Streptococcus mutans* and *Lactobacillus*).

MATERIALS AND METHODS

Preparation of honey samples

This study was carried out with natural, untreated and unpasteurized honey samples obtained directly from beekeepers in Hamadan, West of Iran. The samples were originated from blossoms of wild flowers and did not contain artificial preservatives or diluents.

Assay of antibacterial activity

The honey samples were screened for their antibacterial activity, according to the agar well-diffusion method proposed by the CLSI against the two reference strains: (1) *Streptococcus mutans* PTCC 1683 and (2) *Lactobacil-*

lus spp. PTCC 1643. The above bacteria were grown (100 mL) in Trypticase Soy Broth (TSB) at 37°C for 18 hours. The following honey concentrations (w/v) were prepared in sterile saline solution; 5%, 10%, 20%, 50%, and 100% and their antibacterial activity was evaluated against the bacterial strains.

Accordingly, a 100 mL aliquot of each honey dilution was added to each well on blood agar inoculated with bacterial concentration similar to 0.5 Mac Farland tube, and were incubated at 37°C for 18 hours.

Antibacterial activity was assessed by measuring the diameter of the inhibition zones surrounding the wells. Control plates were prepared with no honey added. All assays were repeated 10 times for each honey concentration.

Statistical analysis

Linear regression lines were estimated for determination of the minimum active dilution of each honey against the reference strains. Data were analyzed by non parametrical Chi-Square test and SPSS 12 for Windows (SPSS Inc., Chicago, Ill, USA). The level of confidence was set at 95% ($\alpha=0.05$).

RESULT

The results are shown in Table 1. There was no significant similarity between the two isolates in their sensitivity to honey ($P>0.05$). Honey in concentration more than 20% had significant antibacterial activity on *Streptococcus mutans* and in 100% on *Lactobacillus* ($P<0.05$).

DISCUSSION

Honey inhibits the growth of a wide range of bacterial species in vitro [24]. However, there have been few studies about the efficacy of honey against the two main cariogenic bacteria *Streptococcus mutans* and *Lactobacillus* [25]. Honey is a delicious, natural sweet food and sweeteners are one of the most common causes in developing dental caries.

Thus, the use of a low harmful sweetener in the diet is very important, especially, if it is confirmed that honey has antibacterial activity against cariogenic bacteria in vitro and in vivo. The effective factors in the antimicrobial activity of honey include enzymatic glucose oxidation reaction, physical properties, high osmotic pressure, low water activity, low PH, acidic environment, low protein content, high carbon

Table 1. Antibacterial Effect of Honey on Bacterial Growth

Honey Concentrations	S.M Sample Number	S.M Growth Number (Percent)	L.B Sample Number	L.B growth Number (Percent)
0%(Control)	10	10(100) ^a	10	10(100) ^a
5%	10	10(100) ^a	10	10(100) ^a
10%	10	10(100) ^a	10	10(100) ^a
20%	10	0(0) ^b	10	10(100) ^a
50%	10	0(0) ^b	10	10(100) ^a
100%	10	0(0) ^b	10	0(0) ^b

Superscripts indicate statistically significant difference between values a and b ($P=0.000$).

to nitrogen ratio, low redox potential due to the high content of reducing sugars, a viscosity that limits dissolved oxygen and other chemical agents, phytochemicals, antioxidants, volatiles, flavonoid, beeswax, nectar pollen and propolis [7,25-27]. This study tested honey, something that is used by people in this region and that is available for them. Because the antibacterial activity of honey varies with the source and processing, results of this study showed that natural honey of this region had an antibacterial effect on cariogenic bacteria (*Streptococcus mutans* and *Lactobacillus*) in experimental studies. Other studies reported that the antibacterial activity of honey increased when the honey was diluted [24]. In the present study, results were not in agreement with this finding and 100% honey concentration had an antibacterial effect on two cariogenic bacteria. Honey in dilutions greater than 20% was effective against *Streptococcus mutans*.

These differences may be due to the type of honey that was used, water content of this honey and honey sources. Researchers evaluated honey from different regions in Egypt for antimicrobial effects. While all samples had antibacterial activity on negative and positive gram bacteria, they were ineffective against *Candida albicans* and *Aspergillus niger* [28]. Research showed that *Helicobacter pylori* is sensitive to honey [29], but honey enhanced the growth of bifidobacteria [30]. *Staphylococcus aureus*, the principal contaminant agent in many clinical infections, has developed resistance against several antibiotics and honey is a costly treatment option in this case. Manuka honey has been reported to be highly effective against various pathogens, including methicillin-resistant *Staphylococcus aureus* (mRSA) and vancomycin-resistant *Enterococci* (VRE) [25].

Findings of a study showed that coagulase-negative *Staphylococci*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Enterococcus* species are susceptible to honey [2]. One study

indicated growth inhibition and acid production of *Streptococcus mitis*, *Streptococcus sorbinus* and *Lactobacillus casei* with 7%, 7.5-8.5% and 8-12% of honey [1].

It was reported that the antibacterial activity of wild flower honey significantly decreased *Streptococcus mutans* in the head and neck irradiated cancer patients that had xerostomia due to radiation [31]. Results of this study were in agreement with the findings of previous studies and demonstrated that honey had antibacterial activity (*Streptococcus mutans* and *Lactobacillus*). Another result which was obtained in this study, was the antibacterial effect of honey tested in concentrations higher than 20% for *Streptococcus mutans*. However, the greatest inhibition of two the bacteria was seen at 100% concentration. A study exhibited that other honeys such as manuka had good antibacterial activity at very low concentrations [21].

An explanation for the difference of results may be due to methodological difference between the studies because the agar dilution method used by others were slightly different from that used in this study. However, it can be due to variation in the composition of the honey being used [21].

In addition, the other honeys have no similar effect on all the bacteria. Although the data obtained from this study could not fully represent the profile of the antibacterial activity of honey in cariogenic bacteria and confirm the use of honey clinically, the present study established that honey had in vitro antibacterial activity and honey might be useful for the prevention of dental caries. Further studies are recommended regarding the antibacterial activity of honey in clinical situations.

CONCLUSION

Under limitations of this study, results demonstrated that natural honey had an antibacterial activity on *Streptococcus mutans* and *Lactobacillus* bacteria. This effect dependent on to the concentration of honey used.

REFERENCES

- 1- Molan PC. The potential of honey to promote oral wellness. *Gen Dent*. 2001 Nov-Dec;49(6):584-9.
- 2-Maeda Y, Loughrey A, Earle JA, Millar BC, Rao JR, Kearns A et al. Antibacterial activity of honey against community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA). *Complement Ther Clinl Pract*. 2008 May;14(2):77-82.
- 3-Zuma A, Lulat A. Honey—a remedy rediscovered. *J R Soc Med*. 1989 Jul;82(7):384-5.
- 4- al Somal N, Coley KE, Molan PC, Hancock BM. Susceptibility of *Helicobacter pylori* to the antibacterial activity of manuka honey. *J R Soc Med*. 1994 Jan;87(1):9-12.
- 5- Ali AY. *The Holy Quran: Translation and Commentary*. 2nd ed. USA: American Trust Puplications,1977.
- 6- Holy Quran, Sure Nahl, The Bees, Aye 68-9.
- 7- Al-Mamary M, Al-Meerri A, Al-Habori M. Antioxidant activities and total phenolics of different types of honey. *Nutr Res*. 2002;22:1041-17.
- 8- Cook NC, Sammon S. Flavonoids- chemistry, metabolism, cardioprotective effects, and dietary sources. *J Nutr Biochem*. 1996;7:66-76.
- 9- Lusby PE, Coombes A, Wilkinson JM. Honey: a potent agent for wound healing. *J Wound Ostomy Continence Nurs*. 2002 Nov;29(6):295-300.
- 10- Ladas SD, Haritos, DN, Raptis SA. Honey may have a laxative effect on normal subjects because of incomplete fructose absorption. *Am J Clin Nutr*. 1995 Dec;62(6):1212-5.
- 11-Ali AT, Chowdhury MN, Al-Humayyd MS. Inhibitory effect of natural honey on *Helicobacter pylori*. *Trop Gastroenterol*. 1991 Jul-Sep;12(3):139-43.
- 12- Ali AT. Prevention of ethanol-induced gastric lesions in rats by natural honey, and its possible mechanism of action. *Scand J Gastroenterol*. 1991 Mar;26(3):281-8.
- 13- Mobarok Ali AT. Natural honey exerts its protective effects against ethanol-induced gastric lesions in rats by preventing depletion of glandular nonprotein sulfhydryls. *Trop Gastroenterol*. 1995 Jan-Mar;16(1):18-26.
- 14-Subrahmanyman M. A prospective randomized clinical and histological study of superficial burn wound healing with honey and silver sulfadiazine. *Burns*. 1997 Mar;24(2):157-61.
- 15-Efem SE, Udoh KT, Lwara CI. The antimicrobial spectrum of honey and its clinical significance. *Infection*. 1992 Jul-Aug;20(4):228-9.
- 16- Molan PC. Honey for treatment of infections. *The New Zealand Beekeeper* 216: 19-20.
- 17- Munstedt K, Lang U. Honey's wound healing properties. *Am Bee J*. 1997;137(4):296-97.
- 18- Iurlina MO, Fritz R. Characterization of microorganisms in Argentinean honeys from different sources. *Int J Food Microbiol*. 2005 Dec;105:297-304.
- 19-Boukraa L, Amara K. Synergistic effect of starch on antibacterial activity of honey. *J Med Food*. 2008;11(1):195-8.
- 20- Molan PC. The antimicrobial activity of honey: 1. The nature of the antibacterial activity. *Bee World*. 1992;73(1):5-28.
- 21- Lusby PE, Coombes AL, Wilkinson JM. Bactericidal activity of different honeys against pathogenic bacteria. *Arch Med Res*. 2005 Sep-Oct;36(5):464-7.
- 22- Alnaqdy A, Ai-Jabri A, Al Mahrooqi Z, Nzeako B, Nsanze H. Inhibition effect of honey on the adherence of *Salmonella* to intestinal epithelial cells in vitro. *Int J Food Microbiol*. 2005 Sep;103(3):347-51.
- 23- Miorin PL, Levy Junior NC, Custodio AR, Bretz WA, Marcucci MC. Antibacterial activity of honey and propolis from *Apis mellifera* and *Tetragonisca angustula* against *Staphylococcus aureus*. *J Appl Microbiol*. 2003;95(5):913-20.

- 24- Mandal MD, Mandal S. Honey: its medicinal property and antibacterial activity. *Asia Pac J Trop Biomed* 2011;154-160
- 25-Lin SM, Molan PC, Cursons RT. The in vitro susceptibility of *Campylobacter* spp. to the antibacterial effect of manuka honey. *Eur J Clin Microbiol Infect Dis.* 2009 Apr;28(4):339-44.
- 26- Irish J, Carter DA, Blair SE, Heard TA. Antibacterial activity of honey from the Australian stingless bee *Trigona carbonaria*. *Inter J Antimicrob Agents.* 2008 Jul;32(1):89-90.
- 27- Cooper RA, Molan PC, Harding KG. Antibacterial activity of honey against strains of *Staphylococcus aureus* from infected wounds. *J R Soc Med.* 1999 Jun;92(6):283-5.
- 28- Roderick JW, Lise KB, Yinrong Lu. Identification and quantitative level of antibacterial components of some New Zealand honeys. *Food Chemistry.* 2000;70:427-35.
- 29- Sato T, Miyata G. The nutraceutical benefit, part III: honey. *Nutrition.* 2000 Jun;16(6):468-69.
- 30- Chick H, Shin HS, Ustunol Z. Growth and acid production by lactic acid bacteria and bifidobacteria in skim milk containing honey. *J Food Sci.* 2001;66:478-81.
- 31 – Sela M, Maroz D, Gedalia I. *Streptococcus mutans* in saliva of normal subjects and neck and head irradiated cancer subjects after consumption of honey. *J Oral Rehabil.* 2000 Mar;27(3):269-70.